


Amendment to the claims:

1. (Currently Amended) A cancer-specific radiolabeled conjugate of the formula:




(I)


wherein B\* represents ~~uracil~~thymidine substituted with a radionuclide; R represents H, OH, or O-L-DHT, L being a cleavable bifunctional linking moiety and DHT is 4-dihydrotestosterone which is bound through its hydroxyl substituent to said linking moiety; and R' represents ~~H or a phospho group or a substituted phospho group~~ having the formula - PO(OR<sub>a</sub>)(OR<sub>b</sub>), -PO(OR<sub>a</sub>)(ODHT) or -PO(ODHT)<sub>2</sub>, R<sub>a</sub> and R<sub>b</sub> being the same or different and representing H or lower alkyl having 1-8 carbon atoms and DHT is as previously defined, with the proviso that at least one of the R and R' substituents comprises a DHT moiety.

2. (Original) The conjugate of claim 1, wherein said radionuclide is an Auger electron-emitting radioisotope selected from the group of <sup>123</sup>I, <sup>125</sup>I, <sup>77</sup>Br, <sup>80m</sup>Br, <sup>195m</sup>Hg and <sup>113</sup>Sn.
3. (Original) The conjugate of claim 1, wherein said radionuclide is selected from the group of <sup>131</sup>I, <sup>32</sup>P, and <sup>225</sup>At.
4. (Original) The conjugate of claim 1, wherein said radionuclide is <sup>125</sup>I.


5. (Original) The conjugate of claim 1, having the formula:




6. (Original) The conjugate of claim 1 having the formula:




7. (Original) The conjugate of claim 1 having the formula:



8. (Original) The conjugate of claim 1 having the formula:



9. (Original) The conjugate of claim 1 having the formula:



10. (Original) A method of treatment of a malignant tumor in a patient in need of such treatment, said tumor comprising cells having androgen receptor, said method comprising administering a therapeutically effective amount of the radiolabeled conjugate of claim 1.
11. (Original) The method of claim 10, wherein said cells are prostate cancer cells.
12. (Original) The method of claim 10, wherein said cells are ovarian cancer cells.
13. (Original) The method of claim 10, wherein said cells are breast cancer cells.
14. (Original) The method of claim 10, wherein said radiolabeled conjugate is administered by a method selected from the group consisting of intravenously, intraperitoneally, and intratumor.
15. (Original) The method of claim 14, wherein said method of administration is administered intravenously.

16. (Original) The method of claim 14, wherein said conjugate is administered periodically for a term of years.
17. (Original) The method of claim 16, wherein said conjugate is administered daily.
18. (Previously Presented) A method for the diagnosis of a malignant tumor in a patient, said tumor comprising cells having androgen receptor, said method comprising administering an effective amount of a radiolabeled conjugate of claim 1, and then imaging the tumor scintigraphically.